

CO25_COM_POA4_054491





Food Safety Conversation No2: Personal Hygiene

FS.SC.002.03

Key Learning What You Need To Know ☐ Hands must be washed before handling ready to eat foods. Good personal hygiene can help **6 STEPS TO EFFECTIVE HAND WASHING** minimise cross contamination ☐ Hands must also be washed after visiting the toilet, handling rubbish, handling or preparing raw food, and ensure the food you cleaning, smoking or eating. prepare, cook and serve to your ■ Effective Handwashing Technique: customers is safe to eat Wet vour hands with warm water Apply hand soap (recommended Ecolab Epicare 5c) Lather and scrub hands for at least 30 seconds Rinse for 10 seconds Suitable hat or hair Turn off the tap using a paper towel COMPASS covering Use additional paper towel to dry hands. (only sleepe ☐ Hair must be clean, neat and tied back or kept covered if handling open food. ☐ With the exception of plain wedding rings and plain sleeper earrings, jewellery must not be worn in good condition while preparing food. Clean hands ☐ Uniform (if provided) must be clean and put on until you arrive at work. Short nails No lewellery ☐ Wearing gloves has not been proven to be a safer method of handling, when food compared to the Clean use of effective hand washing techniques as cross contamination from raw to high risk food can still protective occur and wearing gloves can give a false sense of security. apron Suitable work ☐ Hand to mouth contact must be avoided while carrying out food handling activities so eating, drinking wear for and smoking must be avoided while on duty. indoor use ☐ All food handlers must report signs of illness immediately to their line manager. All food handlers Suitable nor slip shoes - no suffering from symptoms of nausea, stomach cramps, vomiting or diarrhoea must be excluded from onen toes work and not return until they have been symptom free for 48 hours. **HACCP Stages Colleague Validation More Information** Give examples of when hands must be washed. More information can be found within the Good How long should hands be washed to ensure bacteria are removed effectively? All food handling stages Hygiene Practice Guide No: 2 Personal What type of jewellery is permitted? Hygiene and the HSE website What time period must food handlers remain symptom free before returning to work?

CO22_COM_POA4_030445.indd 3 03/03/2022 15:49





Food Safety Conversation No3: Pest Control

FS.SC.003.04

Key Learning			What You Need To Know	
Following this conversation, you will be able to effectively identify signs of pest activity so that you can take appropriate action	 spilled product Do not leave for storage area in Make sure are Check your wo packaging, spi 	t etc. ood deliveries outside or unattended - mmediately; do not leave deliveries ou as are kept clean and tidy, and that wa	aste is removed regularly. as contaminated food, droppings, chewed	
	Area	Signs of Activity	Corrective Action	
Post Control C	Floor	Look for pests (live or dead) and signs of pests including droppings, urine stains, and grease or gnaw marks. Look along skirting boards and under equipment.	Dispose of any dead pests. Sweep up droppings, clear urine or grease stains with disposable paper towel. Then mop and sanitise the area.	
	Shelving & Work Surfaces	Look for pests (live or dead), and signs of pests including droppings, urine stains, and grease or gnaw marks. Remove items and check behind objects.	Dispose of any dead pests. Remove any debris or contamination, then thoroughly clean and sanitise the area.	
	Crockery, glassware, cutlery, utensils etc	Remove items from their containers or shelving. Look inside containers or on shelving for evidence of pests.	Remove any debris or other visible contamination and then wash all items in the dishwasher. Where no dishwasher is available items must be washed with an approved pot-washing detergent with disinfecting properties, at the manufacturers stated dilution and adhering to the stated contact time.	E A
	Food	Visually check food and food containers for gnaw marks and holes in packaging and for droppings or pests (live or dead). Look for spilled product on shelving.	Any contaminated packaging must be disposed of.	
	Disposable Packaging	Check that food packaging is intact without evidence of pests, such as gnaw marks and droppings.	Any contaminated packaging must be disposed of.	
HACCP Stages		Colleague Va	alidation	More Information
ReceiptStoragePreparationCookingService	2. Why is good ho 3. What are some	signs of pest damage you should look fousekeeping important for pest controle signs of pest activity? Ou do if you find evidence of pest activ	More information can be found in the Goo Hygiene Practice Guide No: 3 Pest Cont e and the HSE website	

868763_A4_FoodSafetyTable_03_PestControl.indd 1 03/04/2025 09:58





Food Safety Conversation No4: Cross-Contamination

FS.SC.004.03

Key Learning What You Need To Know · Cross-contamination is the transfer of harmful bacteria from raw foods to ready-to-eat foods. Cross-contamination is one of Cross-contamination must be avoided at all times, as ready-to-eat food will not be cooked / re-heated the main ways food poisoning PREVENT CROSS further which would kill any harmful bacteria that have transferred to the food. can occur. In order to look after CONTAMINATION Cross-contamination can be direct, i.e. from one food directly to another, or indirect, where contamination the safety of our customers USE CORRECT COLOUR CODED CHOPPING BOARDS & KNIVES happens via equipment, hands, surfaces, cloths etc. and provide safe food to eat **RAW MEAT** vou need to ensure that cross-**RAW FISH** contamination of food is avoided. You can prevent cross-contamination by: **COOKED MEATS** Checking deliveries to confirm that raw foods are segregated from ready-to-eat foods in the delivery vehicle. **SALADS & FRUITS** Always storing raw foods separate from ready-to-eat foods, either in separate fridges or by storing raw VEGETABLES foods at the bottom of refrigeration units. **DAIRY PRODUCTS** Preparing raw and ready-to-eat foods separately, ideally in separate, designated areas, or by preparing ready-to-eat foods before raw foods and thoroughly cleaning down and sanitising all work surfaces after the preparation of raw foods. Using separate equipment and utensils for the preparation of raw and ready-to-eat foods, and ensuring preparation area they are cleaned and sanitised between use. Designating complex catering equipment such as mincers, slicers, blenders and vacuum pack machines for Ready-to-eat raw and ready-to-eat foods. food preparation Washing your hands between tasks and avoiding touching other work surfaces, equipment or clothing. area Not washing raw meat and poultry as this results in bacteria being splashed onto surrounding surfaces. This work bench can be Using tongs, other utensils or non-touch methods to avoid direct hand contact with food. used for both RAW and **READY-TO-EAT Products** Following colour-coding for chopping boards and any other equipment where appropriate. Raw products will be prepped first and then a co Replacing or resurfacing worn or damaged chopping-boards. **HACCP Stages Colleague Validation More Information** 1. Can you describe the different types of cross-contamination? Receipt More information can be found Storage 2. Give examples of how you would avoid cross-contamination during storage? in the **Good Hygiene Practice** Preparation **Guide No: 4 Cross-Contamination** 3. Give examples of how you would avoid cross-contamination during food preparation?

4. Can you identify the uses of the different colour coded chopping boards?

Cooking

Service

and the HSE website





Food Safety Conversation No5: Food Labelling & Shelf Life

FS.SC.005.02

Key Learning


It is a legal requirement to ensure food is properly labelled with certain information to help you and the customer know what is in the food and how long it is safe to use or consume the food before it must be discarded

COMPASS



- ☐ 'Use By' dates relate to food safety and food must not be cooked, sold, frozen, consumed or used in any way past this date. It is illegal to sell any food that has passed its 'Use By' date.
- □ 'Best Before' dates relate to food quality and food is safe to eat after this date, depending upon the taste, texture and general quality of foods. Products with a 'Best Before' date may be used beyond this date, unless client site rules require 'Best Before' dates to be stringently observed.
- ☐ Do not use ANY out of date foodstuffs past their 'Use By' date.
- ☐ Sandwiches/wraps etc Apply a shelf life of two days (day of production +1).
- ☐ Chilled / Ready To Eat food Wrap or store in sealed containers and date code with a shelf life of 72 hours
- ☐ Foods prepared and cooked in the unit then chilled and frozen have a food label with up to 6 months shelf life.
- ☐ Food Delivery Checks must include a check of product labels to ensure they have sufficient shelf life to fit in with your menu schedule.
- □ All ready to eat foods with 'Use By' dates are checked daily at end of service and logged on the Closing HSE Checklist.
- ☐ Whenever transferring shelf life dates or other product information from the outer packaging or the food manufacturer's label, only the approved Compass food storage labels must be used





HACCP Stages

- Receipt
- Storage
- Preparation
- Cooling & Re-heating
- Food Service & Display
 - Vending

Colleague Validation

- Describe what a 'Use By' date is.
- 2. Describe what a Best Before date is.
- 3. What shelf life would you give in unit made sandwiches?
- . What shelf life would you give in unit made frozen food?

More Information

More information can be found within the Good
Hygiene Practice Guide No: 5 Food
Labelling & Shelf Life and the HSE website

CO22_COM_POA4_030445.indd 6 03/03/2022 15:49





Food Safety Conversation No6: Food Delivery & Storage


FS.SC.006.03

Key Learning

What You Need To Know

Properly inspecting and checking delivered food products is an essential step to ensure food quality and safety. Storing food under the right environmental conditions protects it from contamination and prevents bacterial growth.

- ALWAYS ensure raw and ready-to-eat foods are kept separate during delivery, receipt and storage.
- Check all food deliveries for shelf life, damaged packaging, and signs of pest damage or infestation. Record all checks on the Food Delivery Record form.
- Check the temperatures off all chilled and frozen food deliveries. Ensure the foods are within the required temperature ranges. Frozen foods must be -18°C to 23°C and chilled foods must be at or below +5°C.
- For all chilled or frozen deliveries either take a copy of the printed vehicle digital temperature
 display reading, take between pack temperature readings using a dedicated probe thermometer
 or use an infrared thermometer to check the temperature of foods. In ROI a physical between pack
 temperature reading is required.



- Ambient foods must be stored in cool, dry and well-ventilated conditions.
- Make sure all foods are labeled and dated, and adopt the "first in first out" rule when replenishing stock.
- Raw foods must be separated from cooked and ready-to-eat foods, ideally by storing them in separate fridges. Alternatively store ready-to-eat foods above eggs and unwashed fresh produce.
 Raw meat, poultry and fish must be stored below all other foods.
- Dry goods must be stored off the floor on suitable shelving, with access all round for cleaning and inspection. Opened packages must be sealed or decanted into suitable lidded containers.
- Fridge temperatures must be checked and recorded twice a day, freezers once per day. The first check should take place within 1 hour of the kitchen opening.
- A food simulant (lard, jelly, water bottle etc) must be used for manual fridge temperature checks, either using a probe thermometer or an infrared thermometer.
- Check the accuracy of all probe and infrared thermometers every month.









HACCP Stages

Colleague Validation

More Information

ReceiptStorage

- 1. What should you check during food deliveries?
- 2. What are the correct operating temperatures for refrigerators and freezers?
- 3. How do you avoid cross contamination of ready-to-eat foods during storage?
- 4. How do you check the operating temperature of refrigerators?

More information can be found within the **Good Hygiene Practice Guide No: 6 Food Delivery & Storage** and the HSE website

859101_A4_FoodDelivery_Storage_table.indd 1 28/01/2025 15:26





Food Safety Conversation No7: Food Preparation

FS.SC.007.02

Key Learning What You Need To Know ☐ There are principally 4 sources of food contamination: The preparation stage of any food operation is the most Physical important with respect to Chemical controlling the risks of Bacterial Allergenic contamination. It is essential in our role as a food handler to ☐ Always wash hands in warm water with soap and dry them using disposable paper towels before handling any food. know these risks and adopt the control measures to minimise ☐ Follow any food safety instructions on food packaging regarding the preparation and handling of food. food contamination ☐ Prepare food as close to service time as possible, keeping perishable foods under refrigeration. ☐ Where possible provide separate work areas for raw and ready to eat foods. ☐ Where this is not possible segregate via time and thoroughly clean and sanitise areas between use. ☐ Follow your units colour coded chopping board system if one is implemented. ☐ Do not use complex equipment for both raw and ready-to-eat foods e.g. vacuum packers, food slicers. food mixers and food processers. ☐ Thoroughly wash salad and fruit to be sold as ready to eat with clean water to remove visible dirt. ☐ Ensure food allergens are handled and prepared carefully to avoid cross contamination, use separate boards and utensils. ☐ Make sure you know what ingredients are in a food item prepared and made on site in order that the correct allergen information can be given to the customer if requested. ☐ Sanitisers used to clean work surfaces must meet the standard BSEN1276 and be used in accordance to the manufacture's instructions, including the correct dilution and contact times. ☐ Where possible avoid using glass in food handling areas and always check any glassware used for signs of chipping / cracks / breaks and do not use if any damage is found. **More Information HACCP Stages Colleague Validation** 1. What are the different types of food contamination and give examples? 2. How can you avoid cross contamination? More information can be found within the Good Preparation 3. Why do you need to wash fruit & vegetables before use? **Hygiene Practice Guide No: 7 Food** Food Service & Display 4. Why is it important to know what ingredients go into a dish you are preparing? Preparation and the HSE website What checks should you do if using glass products within your kitchen?

CO22_COM_POA4_030445.indd 8 03/03/2022 15:49





Food Safety Conversation No8: Defrosting

FS.SC.008.02

Key Learning If food is not thoroughly

defrosted before cooking, heat will be used to thaw the food rather than cook it. There is a danger that a core temperature of +75°C, which is required to kill any harmful bacteria present, will not be achieved



What You Need To Know

- ☐ Freezing food does not kill harmful bacteria but it will prevent growth. As the food begins to warm during thawing, bacteria begin to multiply slowly
- ☐ All food must be completely defrosted prior to cooking or reheating, unless the manufacturer's instructions indicate the food is to be cooked from frozen.
- □ Factors affecting defrosting times:
 - Defrosting times are faster if food is frozen and then defrosted in smaller portions.
 - Loose wrapping, which traps an insulating layer of air, increases thawing time.
 - Foil wrap can reduce heat radiation and slows thawing time.
 - Tight packaging, such as vacuum packaging or food wrap, has little effect on thawing times.
 - Food will thaw more quickly at ambient temperature. The main disadvantage is that, at higher temperatures, bacteria can multiply more freely and may reach unacceptable levels.
- ☐ It is recommended that all foods are defrosted slowly in a refrigerator (between 0°C and +5°C), ensuring all raw foods are placed at the bottom of the refrigerator, separated from ready to eat foods to avoid cross contamination
- ☐ Place defrosting raw food items in a container or receptacle to ensure meltwater does not cross contaminate other foods
- Only defrost foods in a microwave if recommended by the manufacturers instructions using the defrost setting only. Ensure the food is spread evenly on the turntable and stir frequently.
- Defrosting food at room temperature is permissible if it is defrosted in a controlled environment which is free from sources of contamination. Do not use running cold water to speed up the process.
- ☐ Cook and / or serve any defrosted within its remaining shelf life
- Do not re-freeze any defrosted foods







HACCP Stages

- Preparation
- Cooling & Re-heating
- Food Service & Display

Colleague Validation

- 1. What are the dangers of not thoroughly defrosting food before cooking it?
- 2. What factors can affect the time in which foods are defrosted?
- 3. How would you safely defrost food in a refrigerator?
- 4. How would you ensure food is properly defrosted in a microwave?
- Can you re-freeze defrosted food?

More Information

More information can be found within the Good Hygiene Practice Guide No: 8 Defrosting and the HSE website

CO22 COM POA4 030445.indd 9 03/03/2022 15:49





Food Safety Conversation No9: Cooking

FS.SC.009.03

Key Learning What You Need To Know Cooking is a critical step to ☐ Food poisoning micro-organisms are killed through the correct Time and Temperature combination. ensure that any bacteria that ☐ Food must be cooked thoroughly to achieve a core temperature of +75°C. may be present in food are ☐ Temperature check all protein foods using a sanitised probe thermometer to ensure completion of the completely killed and the food is cooking process, and record the temperature on the Food Production Temperature record form or the safe to eat for your customers digital HACCP equivalent. □ Never undercook rolled joints / minced or diced meat / poultry / pork as these are high risk food items and require thorough cooking to reduce the risk of food poisoning bacteria being present. □ Some whole cuts of meat and some fish products may be cooked to a lower temperature. For whole cuts of meats required "pink" or "rare" check to ensure the outer surface of the meat is fully sealed and browned off before serving. Whole pieces of fish (e.g. tuna steaks) can be served 'rare' as long as they have been fully seared on the outside. ☐ Also visually check food to ensure it has been cooked thoroughly: Check the centre of fish or near the bone to make sure that the colour and texture has changed. Shellfish such as prawns and scallops will change in colour and texture when they are The juices from pork, rolled joints and poultry should be clear and not have any red of pink in them. Ensure poultry and minced meat products are not pink inside Check that pizza bases are cooked and that the toppings are bubbling ☐ Food temperature probes must be accuracy checked every month using melting ice or boiling water and recorded in the HSE Logbook or on the Probe Thermometer Accuracy Record Form. **HACCP Stages Colleague Validation More Information** 1. What is the standard Compass cook temperature to ensure thorough cooking? More information can be found in the Good Preparation 2. Give examples of visual checks you can undertake to check food has been cooked properly. Cooking Hygiene Practice Guide No: 9 Cooking and How do you ensure a food probe thermometer is working correctly? Food Service & Display the HSE website

CO22_COM_POA4_030445.indd 10 29/01/2025 15:49





Food Safety Conversation No10: Cooling & Reheating


FS.SC.010.02

Key Learning

Effective and swift cooling techniques allows the safe storage of in unit made foods. Re-heating foods to the correct temperatures ensures any remaining bacteria present in food will be killed through heat



- ☐ Micro-organisms that cause illness and food spoilage can grow between 5°C and 63°C. This range is known as the Danger Zone, and keeping foods out of this zone reduces the risks of food poisoning.
- ☐ Cool food as quickly as possible after cooking and place into refrigerated storage.
- ☐ Cooling times can be reduced by:
 - Portioning the food into smaller containers
 - Cutting or slicing larger joints before cooling
 - Using shallow / pre cooled containers (5cm depth)
 - Using ice baths to rapidly cool the food
 - Rinsing under cold potable water (e.g. for rice or pasta)
- ☐ When cooling food in a blast chiller ensure the chilling process starts within 30 mins of the food being cooked, and that if is cooled to a temperature of 5°C or less before placing into refrigerated storage.
- ☐ When cooling food at room temperature choose a cooler area of the kitchen. Cover the food and leave at ambient room temp for a maximum of 90 mins before placing into refrigerated storage.
- ☐ Record the following information on the Food Production Temperature Record form:
 - Type of food
 - Time & Temperature of the food when cooling began
 - Time & Temperature of the food when transferred to refrigeration
- ☐ If food is cooling slower than expected, break the food down into smaller quantities, transfer food to newly cooled containers or place food container in cold water and ice several times.
- ☐ It is important to re-heat food thoroughly to ensure any remaining harmful bacteria are killed off.
- ☐ Reheat foods to the correct temperature (75°C England / Wales / N.I. 82°C Scotland).
- ☐ Record reheat temperatures on the Food Production Temperature Record form.



Reheating Food



Scotland



HACCP Stages

1.

Preparation

- Cooling & Re-heating
- Food Service & Display

Colleague Validation

- What is temperature range is known as the "Danger Zone" which you must avoid keeping food at?
- 2. What practical steps could you take to help cool food quickly?
- 3. What is the maximum time period allowed to cool food at ambient room temperature?
 - What are the required reheating temperatures?

More Information

More information can be found within the Good Hygiene Practice Guide No: 10 Cooling & Reheating and the HSE website

CO22_COM_POA4_030445.indd 11 03/03/2022 15:49





Food Safety Conversation No11: Food Service & Display

FS.SC.011.03

Key Learning	What You Need To Know	
Displaying food in a safe environment, free from potential sources of contamination will ensure we look after our	Protect food on display by: ☐ Ensuring suitable precautions are taken to minimise the risk of allergen cross-contact on salad bars, deli counters, buffets etc by providing separate service utensils and equipment for different food types. ☐ Labelling all displayed food so that it can be easily identified. ☐ Using sneeze guards or covers where possible to protect displayed food from contamination by customers.	Hold at or above +63°C
customers and serve good quality, safe food.	 □ Washing any raw fruit, vegetables, herbs or other items added to the display for visual effect. Potted herbs must not be placed in display units. Cold Food Display □ Ready-to-eat foods must be kept at or below +8°C (+5°C in Rol). □ Ensure cold display units are operating to temperature before loading them. 	Food Limit to 2hrs if displayed at ambient
Food Service & Display	 □ Do not display food above the load line of a chilled food well and allow space in-between food containers to allow cold air to circulate effectively. □ Do not use items such as wooden crates or boards, bricks, matting or wicker baskets to elevate food as this prevents the transfer of cold air to the food. □ Cold food may be displayed out of chilled storage (above +8°C / +5°C in Rol) for a single period of up to 4 hours only. After this period, it must be disposed of. □ Deli Bars 	Hold at or below +8°C (5°C in Ireland)
	☐ Food must be cooked or reheated thoroughly before hot holding begins. ☐ Food in hot holding must be kept at or above +63°C. ☐ Hot food may be displayed out of temperature control for a single period only of up to 2 hours post cooking/reheating. ☐ Preheat hot holding equipment e.g. bain-maries/hot cabinets before you put any food in it. Monitoring	Limit to 4hrs if displayed at ambient
Section Sec	 □ Monitor the temperature of one protein food item on each cold / hot display counter and where relevant, one rice dish, and record it on the Food Service Temperature Record from. In Rol all protein foods should be monitored. □ In cold display units a simulant should be used to monitor temperatures. □ Take temperature readings at the beginning of each service period and every 90 minutes for hot food and every 3 ½ hours for cold food. □ Where food is stored outside of temperature control, a documented system must be in place to ensure the permitted time period is not exceeded. 	Mixed Limit to 2hrs if displayed at ambient
HACCP Stages	Colleague Validation	More Information
Hot Hold Food Service & Display	 What temperature should hot food on display be held at? How frequently should you monitor and record the temperature of hot and cold food on display? What are the exemption times for holding hot and cold food out of temperature control? How long should a mixed buffet (hot & cold foods) be left out for clients? 	More information can be found in the Good Hygiene Practice Guide No: 11 Food Service & Display and the HSE website

CO25 COM POA4 053906





Food Safety Conversation No12: Cleaning

FS.SC.012.02

Key Learning What You Need To Know **Definitions** Cleaning food premises and equipment ensures that harmful ☐ Cleaning – The act of removing dirt using water and a cleaning agent ☐ Disinfection – To clean using a substance that kills food poisoning micro-organisms food poisoning bacteria cannot ☐ Sanitising – A cleaning process of reducing microbiological contamination to a level that is acceptable multiply and cause harm, as well □ 2 stage clean – A cleaning program that involves both general cleaning and disinfection / sanitising as helping to ensure the ☐ Clean as you go – Removal of general debris / cleaning up spillages, throughout the task premises are kept clean, tidy ☐ Deep Clean – The periodic cleaning of walls / floors / ceilings / equipment / ventilation etc which does and pest free not form part of the daily cleaning schedule Guidance Stage 1 ☐ Use disposable cloths or paper towels where possible for cleaning food surfaces. COMPASS using correct concentration as ecommended by manufacture ☐ Where separate areas for raw and ready to eat foods have Remove food debris with pape towel or suitable tool been provided, avoid cross contamination by using colour coded cleaning equipment such as cloths and designating cleaning spray Surface clean with a colour coded / disposable cloth or bottles specifically for use in that area. ■ A two stage cleaning process will be needed where food contact surfaces, sinks or equipment have come into contact with raw meat Stage 2 or unwashed fruit and veg and always at the end of the day. · Re-spray sanitiser ☐ Store cleaning equipment and chemicals away from food in a suitable Wipe with disposable cloth or cleaning store to avoid contamination of food blue paper towel and leave in contact with work surface for ☐ Re-usable cloths and towels should be effectively segregated (e.g. colour-coded) and suitably washed at high temperatures at the end of each day in order to destroy bacteria especially E.coli O157 **HACCP Stages Colleague Validation More Information** What is the difference between disinfection and sanitising 1. More information can be found within the 2. Describe a 2 stage cleaning process All Stages **Good Hygiene Practice Guide No: 12** 3. How would you avoid cross contamination when cleaning Cleaning and the HSE website

CO22_COM_POA4_030445.indd 13 03/03/2022 15:49

What would you do to demonstrate "clean as you go" within your work area





Food Safety Conversation No13: Allergens

FS.SC.013.03

Key Learning

Understanding and following the allergen procedures will ensure the safety of your customers and potentially avoid serious injury or death due to allergic reactions

There are 14 major food allergens which must be declared. However. people can be allergic to any food or ingredient, such as pea protein, stone fruit or coconut.









What You Need To Know



















Retter	Riges													
	0	0	0	0	0	0	0	0	0	0	9	0	0	0
	time	Creature.	Commen	Tarr	100	Same	-	Street,	Mutuell	Femili	Title See	Dame	better	Name of
Secretary	_	_	_	_	_	_	_	_		_	-	_	_	-
compagnio de de		d'anne		_						w	_		1	-
Service .		100		1			*							
Station.				7										
Sant Sant														
-														
had been been been														
differ from .														
		d'intest								MC				
Frencher Spc		MIC BOOK												- 4
-		100												
to taribone														
departure .				1										
		-												



- ☐ A food allergy happens when the immune system triggers a reaction to a food or ingredient (allergen), causing symptoms such as rashes, swelling, breathing difficulty and anaphylaxis, Symptoms usually develop very quickly and can be fatal.
- ☐ A food intolerance is an adverse reaction to food or ingredient which doesn't involve the immune system. Symptoms include stomach upsets, bloating and headaches and usually develop more slowly.

Avoiding Allergen Cross-Contact

- ☐ Use separate chopping boards and utensils for each dish.
- ☐ Clean down work surfaces, equipment and your hands between handling know allergen containing foods and before preparing an allergen free meal.
- ☐ Keep opening dry goods in sealed containers or tightly covered.
- ☐ Use separate fryers for known allergen foods (e.g. scampi / battered fish) and non allergen foods (e.g. chips)
- ☐ Use separate cooking equipment / utensils when cooking an known allergen free item

Kitchen Process

Menu planned on the Source











older printed

HACCP Stages

- Food Storage
- **Cross Contamination**
- Preparation
- Cooking
- Service
- Hospitality

Colleague Validation

- 1. Explain the differences between an allergic reaction and an intolerance?
- 2. What are the 14 major food allergens?
- 3. Why is it important to follow a recipe without making any changes?
- 4. Give examples of how can you avoid allergen cross-contact?
- 5. How do we inform customers of what allergen ingredients are in a menu item?

More Information

More information can be found within the Good Hygiene Practice Guide No: 13 Allergens and the HSE website

CO22 COM POA4 030445.indd 29/01/2025 15:49





Food Safety Conversation No14: Catering Premises & Equipment

Key Learning

What You Need To Know

Good design, construction, location of equipment and workflow of a kitchen help minimise the risk of crosscontamination and aid cleaning and maintenance.

Structural Requirements

- Kitchen design should allow workflow from delivery, to preparation, through to cooking and service.
- Separate storage should be provided for raw and ready-to-eat foods.
- Walls, floors and work surfaces should be constructed of materials suitable to allow effective cleaning.
- Food premises should be proofed to prevent pest access and harbourage.
- Suitable mechanical or natural ventilation must be provided to ensure that heat and/or humidity does not build up.
- · Wash hand basins, supplied with hot and cold water, bactericidal soap and single use towels, must be provided and used only to wash hands.
- Separate sinks designated for food and sinks for equipment washing should be available. Where there is only one sink, this must be thoroughly cleaned and sanitised between uses (not acceptable in ROI).



Food Equipment

- · Separate, clearly designated, complex equipment must be provided for raw and for ready-to-eat foods, to avoid cross-contamination.
- Wood or wooden products are not recommended unless they are well maintained and can be effectively. cleaned and disinfected.
- Lighting must be sufficient to allow for safe working and easy inspection of all areas.
- Food service ware and crockery should be checked for damage, such as chips and cracks, before use. Any damaged items must be discarded to prevent foreign body contamination.

Maintenance

- Food contact surfaces, equipment and utensils must be maintained in good condition and checked before use. If damaged they should be removed from service and discarded or clearly labelled "Do Not Use"
- Any defective structure, equipment or utensils should be recorded on the Compass Repair & Maintenance Record form, or client based system, and reported to the correct maintenance provider to fix.







HACCP Stages

All Stages

Colleague Validation

- 1. What should wash hand basins be provided with?
- 2. Where a single sink is provided for equipment and food, what should you do between uses?
- 3. What should you check for before using food equipment?
- 4. Give examples of separate raw and ready-to-eat food equipment?
- 5. How would you report defective food equipment?

More Information

More information can be found within the **Good Hygiene Practice Guide No:14 Catering Premises & Equipment** and the **HSE** website

859101 A4 CateringPremises 14 Equipment table.indd 29/01/2025 10:59





Food Safety Conversation No15: Vac Packing

FS.SC.015.04

Key Learning	What You Need To Know	
Vac Packing machines are a complex piece of equipment and must be kept clean and clearly labelled as either for "raw" or "ready to eat" foods	 Separate vacuum packing machines MUST be used and clearly labelled for "raw" or "ready-to-eat" foods only Vacuum packing bags used for raw or ready-to eat foods must be stored separately and ideally within reach of the designated vacuum packer and be clearly labelled. 	
Vacuum Packing Packing	 Only those listed on the Authorised User List (after completing training) are permitted to use and clean the vacuum packing machines. Close attention should be paid to the hidden areas and removable plates, and the minimum contact time on the sanitiser observed. Good personal hygiene is important at all times, and food handlers MUST wash their hands thoroughly before and after using the vacuum packing machines. Where possible, direct handling of the food should be kept to a minimum. 	READY-TO-EAT USE ONLY O : 3- O DOOLD H
READY-TO-EAT USE ONLY Standard Park May Standard	 Ensure every packet is suitably and sufficiently sealed with a tight fit round the food and the seal intact. Check packets for excess air and/or leaks to minimise the risk of contamination and subsequent growth of bacteria. Vacuum packed ready-to-eat high-risk food must be stored at a maximum of 5°C or below. Date labelling (shelf-life) for in-unit made chilled foods must not exceed 7 days from the date and time of production (3 days in Republic of Ireland) and 6 months (3 in Republic of Ireland) for frozen in unit made foods Clear 'production' and 'use-by' dates should be put on all packets using the Compass date labels. Any out of date chilled and frozen vacuum-packed products must be discarded, even if the food appears 	RAW USE ONLY
	acceptable to the senses. Uacuum packers must be serviced annually by a competent engineer, and records retained on file. Colleague Validation	More Information
All Stages	 Why is there a need to have separate raw and ready-to-eat vacuum packers? Should raw and ready-to-eat vacuum packers and packaging materials be clearly labelled? All food handlers who may use the vacuum packers appropriately trained, and understand the cross-contamination risks and hazards associated with vacuum packers? Are the vacuum packers being adequately cleaned before and after use? 	More information can be found within the Good Hygiene Practice Guide No: 15 Vac Packing and the HSE website

CO22_COM_POA4_030445.indd 16 03/03/2022 15:49





Food Safety Conversation No16: Bar Service

FS.SC.016.02

Key Learning	What You Need To Know	
The quality and shelf life of beer can be affected by wild yeasts, mould and spores, therefore it is important to have a regular cleaning regime in place for structures, equipment and plant.	Beer Line Cleaning: Ensure that full PPE is worn when carrying out the beer line cleaning. Use only the authorised beer line cleaning detergent available from Ecolab. Dilute the detergent with water following the manufacturer guidelines. Draw the detergent solution through at least 3 times during the line cleaning Aim to leave the detergent solution in place on each pull for about 10 minutes. Do not leave for longer than 2 hours as this can taint the pipes. Flush through with a minimum of 8 pints of clean water at the end of line cleaning to remove all traces of detergent. Check with litmus paper that there are no detergent traces. Put signage in place to warn all colleagues that line cleaning is in operation. Leave unused lines charged with water, BUT the lines must be put through the line cleaning process weekly. Ice Machines Ice machines can be a source of Legionella bacteria and E-coli due to unhygienic use and poor cleanliness Ensure that before you proceed to use an ice machine or decant ice that you thoroughly wash your hands first Ice scoops must never be left in the ice machines or ice wells. Ice scoops should be stored in a separate sanitised container and washed in a dishwasher daily	CLEAN Beer Line With beerstone calcium buildup
	 A twice daily clean of the external and hand contact surfaces of the ice machine should be conducted using the sanitiser spray and ensuring a minimum 1 minute contact time. Every two weeks the ice machines should be emptied and cleaned internally and sanitised 	
HACCP Stages	Colleague Validation	More Information
Bar ServiceHospitality	 How many times should you draw the detergent through a beer line before flushing with water? How many pints of clean water should you pull through a beer line after cleaning? What do you do with Ice scoops after use? How frequently do you clean an ice machine? 	More information can be found within the Good Hygiene Practice Guide No: 16 Bar Service and the HSE website

CO22_COM_POA4_030445.indd 17 03/03/2022 15:49





Food Safety Conversation No17: Steamplicity / Esteem

FS.SC.017.02

Key Learning What You Need To Know To understand the specific Steamplicity and Esteem are brand names for a unique product produced by Compass to produce fresh nutritious meals in healthcare, schools and business environments. The method of cooking allows for different requirements for the delivery, products to be cooked together - such as fish and vegetables at the same time. Cooking takes between 3-5 storage, cooking and service of minutes for plated meals 6-10 minutes for bulk meals. Steamplicity plated meals, **Delivery controls:** Steamplicity bulk meals, and ☐ Check vehicle temperature via print out prior to offloading the products Esteem meals ☐ Remove to chilled storage within 30 minutes of delivery ☐ ALL Steamplicity products should be between 0°c and +5°c ☐ If temperature above 5°C destructive product test needs to be taken – refer to cuisine centre for guidance ☐ If between +8°c - +10°c delivery can be accepted if used within 12 hours if not delivery MUST be rejected. ☐ Chill unit must be able to chill below 5°c rapidly and record on the Steamplicity/Esteem delivery record. Storage: ☐ Must be controlled between 0°c- +5°c Plated meals should not be stacked more than two high □ Check stock rotation and shelf life Pick and Pack: ☐ Must be controlled between 0°c- +5°c (can be picked up to +8°C but only for a limited time) ☐ Place in pre chilled transport containers ☐ Late or individual meals to be transported in insulated chilled bags ☐ Record temperature on dispatch pick and pack record – on leaving chilled storage Satellite/ward Storage: ☐ Must be stored between 0°c-5°c and stored separate or above other food items ☐ Remove from chilled storage as close to cooking time as possible Max. 15 minutes Cooking: Check seal is intact and meal in date ☐ Check number on bar coded label and cook on correct programme in microwave with valve facing up. ☐ After cooking leave for 30 seconds for pressure to equalize. ☐ Using cleaned probe check temperature of protein item. Temperature to achieve: Steamplicity plated 82°c / Esteem 82°c / Steamplicity Bulk 75°C ☐ If temperature not achieved return to microwave and boost on 0 for 30 seconds(plated and Esteem / I min for bulk- if temperature still not achieved dispose of the meal. Report fault to manager. ☐ Record each meal temperature on the Steamplicity/esteem cooking record. **HACCP Stages**

	H		14	L	
		000	0000000	- 18	
	F	-			omplet.
	P	4			1
	318				
4					-
3	JA NI		1000	3	

		TY	co	MPASS C	SROUP UI	& IRELANI	0
	57	EAMPI IC	тусы	I ED TE	MPERATI	IRE MONE	TORING SHEET
							aken at least twice daily imorning.
and aftern 10 minute taken by e emperation	oon) 7 da a (chiller i lectronic ire is still ire is abor	rys a week may be or reading, h outside th we 5°C follow	defrost and held and held a set lim ow the h	emperato and take d probe in its, destr (ACCP pl)	are is out; the term the chill sective term in	ide the set. senature ag or or simula ting with a	aren at reast twice daily yearning immits return to the childre within ain. The temperature can be at kept in the childre. If the me at will be required. If the minutes or less.
Worth / Y		iler. U.C. ti	Sean	Unit:	is accept.	icie for au	Area:
Date 1 2 3	Time	Temp	Sig	Time	Temp	Sig	Comments
4 5							
5 7							
8							
10							
12							
16							
16							
18							
20							
22							
24 25							
26 27							
28							
30 31							
-							
Thecked B	W.						Date:

Steamplicity / Esteem sites only

Colleague Validation

- 1. What temperature should Steamplicity/ Esteem products be delivered?
- 2. How would you transfer a late meal and where would you record the temperature.
 - Give the process you would follow to cook a Steamplicity/Esteem meal

More Information

More information can be found within the Good Hygiene Practice Guide No: 17 Steamplicity and the HSE website

CO22 COM POA4 030445 No 17 pg.indd 1 31/03/2022 10:23





Food Safety Conversation No18: Food Waste

FS.SC.021.03

Key Learning What You Need To Know Food waste not only generates Food waste includes spoilage, trimmings, cooking errors, over production and plate waste. dangerous greenhouse gases We have a 3-pronged approach to minimise food waste: when it decomposes but it costs the food industry 1. Prevention - To reduce the amount of food waste that goes unsold/uneaten £2.5b pounds per year. Help 2. Recovery - To repurpose food protect the environment 3. Recycling - To divert food waste from landfill and your units profitability by reducing food waste from The main ways to prevent food waste are: your catering operation. ☐ Tracking Food Waste: recording food waste daily helps understand how much food is being wasted. ☐ Good Menu Planning: regularly review menus to identify slow moving items; take into consideration seasonal trends, the weather, local or national events etc. ☐ Food Storage: ensure food is stored at correct temperatures and in suitable containers, manage stock rotation, and follow manufacturer's instructions when applying shelf-life to opened sauces, condiments, spices etc. ☐ Preparation and Cooking; keep trimmings to a minimum, cook in small batches and cook high value items to order were appropriate. ☐ Food Service and Display: ensure required temperatures are maintained and do not overstock. ☐ Portion control: keep portions consistent and standardise them. ☐ Avoid Overproduction: adjust production to predicted demand and consider reduced service during quieter ☐ Plate Waste: monitor plate waste and adjust menus or portions accordingly. We can repurpose food by using it internally or donating it to our charity partners. Food must be in date and allergen information must be provided. Foods that can be repurposed include packaged ambient foods, fruit and vegetables, pre-packed chilled foods, frozen foods and bakery items. Some hot food can also be repurposed - refer to GHP18 for details. **HACCP Stages Colleague Validation More Information** 1. What are some of the different types of food waste that are generated from kitchens? More information can be found 2. How can we prevent food waste? in the **Good Hygiene Practice** All Stages What types of foods can be repurposed? **Guide No: 18 Food** Waste and the HSE website

883652_A4_FoodSafetyTable_18_FoodWaste.indd 1 11/08/2025 12:41





Food Safety Conversation No19: Food Incidents & Enforcement

ES SC 022 03

		F3.3C.022.02
Key Learning	What You Need To Know	
Dealing with customer food safety	Typical causes of food safety complaints include:	
complaints is important in order to provide good customer service	☐ Allegations of Food Poisoning - Where a customer suspects illness from eating unfit or contaminated food	
and investigate and prevent	☐ Foreign Body Contamination - The physical contamination of food by a foreign body object	
further incidents. Confidently dealing with a food safety	☐ Food Allergen / Intolerance - An allergic reaction or intolerance to a particular type of food consumed	X
enforcement visit will directly	☐ Product Miss-labelling - Where a food item is incorrectly advertised / labelled as something else	
affect the Food Hygiene Rating awarded at the end of the visit	☐ Viruses - Where customers have reported food poisoning type symptoms but may be infected with a food bourn illness, e.g. Norovirus	33
	Dealing with food safety incidents:	
COMPASS Food Safety	☐ Always take the details of any customer complaint, noting their name, contact information, date, time, accurate details of the food eaten and details of any illness or physical injuries sustained.	(**)
Incidents & Enforcement Control House Program Charles for 19	 Report any customer complaint immediately to the Unit Manager or Head Chef and record all customer complaints on the HSE reporting system (AIR2) to ensure appropriate escalation to the Operational Management team and HSE Manager. 	* •
	☐ Undertake a full investigation of all food safety complaints using the appropriate checklist to ensure the route cause can be determined and prevented in future	
	Managing Enforcement Officer visits:	As As
Source Inc. Section 1997 (Section Inc.) 1998 (SAFE) Source Inc. Section 1998 (SAFE) Source Inc. Section 1998 (SAFE) Source Inc. Section 1998 (SAFE)	□ Environmental Health Officers will visit food businesses either routinely to undertake spot checks and issue a Food Hygiene Rating (FHR) or may also visit following a compliant made by a customer regarding an incident.	
	□ EHO's will look at 3 main areas to determine the Food Hygiene Rating – Food Safety Procedures / Structure & Cleanliness / Confidence in Management	
HACCP Stages	Colleague Validation	More Information
Date code & labellingTemperature recordsTraining records	 Describe the typical causes of food safety complaints made within food premises? List the main things you would need to note down when being informed of a food complaint? How would you report food safety incidents? Who would visit your premises to undertake routine food safety inspections? What are the 3 areas of food safety compliance an EHO would look at to determine your Food Hygiene Rating? 	More information can be found within the Good Hygiene Practice Guide No: 19 Food Incidents & Enforcement and the HSE website

CO22_COM_POA4_030445.indd 19 03/03/2022 15:49





Food Safety Conversation No20: Sous Vide

FS.SC.026.01

Key Learning

In sous vide cooking the temperatures used are often much lower than those used for general cooking, and the cooking times much longer, which means that food is in the temperature danger zone for extended periods. As a result, there is an increased risk that food poisoning bacteria can **survive** and **multiply**. Therefore, it is essential that the guidance is followed, and cooking is carried out properly when using sous vide.



What You Need To Know

- □ Sous vide is considered a high-risk activity by the EHO. They will pay particular attention to your process if you use sous vide in your sites.
- ☐ The sous vide GHP 20 must be followed carefully and all associated records to be in place.
- ☐ Senior Kitchen Staff must hold a Level 3 Food Safety certificate or above before embarking on using sous vide in their kitchens.
- ☐ Kitchens must not deviate from the cooking matrix unless specifically given authorisation from the HSE team in writing.
- ☐ Water baths need to be calibrated at least once a year to an accuracy of 0.1'c
- ☐ Water bath front panel LCD temperature display must be calibrated against daily and a correction figure noted if necessary.
- ☐ All items to be cooked must be at room temperature before cooking starts.
- Cooking time only starts when the bath temperature settles to the correct temperature.
- No fish to be cooked sous vide.
- □ No food items being sous vide cooked to be thicker that 5cm or larger than 2kg in weight per bag.





HACCP Stages

- Food preparation
- Cooking

Colleague Validation

- 1. What are the dangers of not following the cooking matrix?
- 2. What factors can affect the length of time in which food cooking takes place?
- 3. How would you best chill sous vide cooked food?
- 4. How would you ensure the water bath is at the correct temperature?
- 5. How long can you store sealed foods once cooled?

More Information

More information can be found within the Good Hygiene Practice Guide No: 20 Souse Vide and the HSE website

CO22_COM_POA4_030445.indd 20 03/03/2022 15:49





Food Safety Conversation No21: Natasha's Law

FS.SC.024.02

Key Learning

The Food Information Regulations (Amendments) 2019, also known as Natasha's Law, has been introduced to ensure customers have access to the full ingredient and allergen information for pre-packed foods to allow them to make an **informed choice** before purchase.



What You Need To Know

□ Pre-Packed Foods for Direct Sale (PPDS) includes foods which are made or prepared in the unit from which it is sold, and it is packaged before being offered for sale. It can be either foods the customer select themselves (e.g. from a display unit), as well as products behind the counter or sold at mobile or temporary outlets.

Foods That Require Labelling

Sandwiches and bakery products which are packed in unit before a consumer selects or orders them

Fast food packed before it is ordered, such as a burger under a hot lamp

Products that are pre-packaged on site ready for the lunchtime sale, such as pizzas, pasties, salads and pasta

Foods packaged and then sold elsewhere on the premises at a mobile / temporary site or vending machine by the same food business

Pre-made drinks placed into lidded containers before being ordered by a customer

Foods That Do Not Require Labelling

Traditional Counter Service – e.g. over the counter meal service

Deli Bar / Salad Bar – Any food to go where customer selects from an open food display and it is packaged after section

Pre-ordered Packaged Primary School Meals where parents have selected the meal

Patient / Resident Meal Service – Any form of hospital or care home patient feeding service whereby the food is ordered ahead of consumption

Hospitality – Any form of food ordered by the client for an event or meeting whereby an informed choice is made

Packed Lunches / Picnics / Hampers – Any pre-ordered food whereby an informed choice is made before the food is packaged

Distance Selling – Any foods purchased or pre-ordered via a website or mobile app

Bought In Food – Any packaged foods bought in from a supplier or CPU will already be labelled

Note: Products which are packaged but sold through a pre-order service (schools / patient feeding / hospitality etc) are excluded from legislation as the 'informed choice' has already been made.

☐ There is a wide range of size and type of labels available depending on the product and complexity of the items. Should be labelled, but aren't must be removed from sale until the items can be labelled.









HACCP Stages

- Food preparation
- Cooking

Colleague Validation

- 1. Should a pre-made sandwich that has been wrapped in clingfilm be labelled?
- 2. Does a pie that is in an open take-away tray need to be labelled?
- 3. Does a pre-made juice in a lidded container require labelling?
- 4. What should you do if you notice that a pre-packed item, such as a packaged salad, doesn't have a label?

More Information

More information can be found within the Good Hygiene Practice Guide No: 21 Natasha's Law and the HSE website

CO22_COM_POA4_030445.indd 21 03/03/2022 15:49





Food Safety Conversation No22: Food Transportation

FS.SC.027.01

Key Learning	What You Need To Know	
B	Typical controls of food transportation include:	
Poor transportation can put our customers at risk, this will help you	☐ Transport cold food at +8°C or lower (5°C in Ireland)	
reduce the risks around microbiological, chemical and	☐ Transport hot food at +63°C or hotter	
physical contamination, and demonstrate proper temperature	☐ High risk foods served cold which are transported above +8°C (5°C in Ireland) must be used / served within 2 hours	Take Est Accord
controls	☐ High risk foods to be served at ambient temperature which is transported above +8°C (5°C in Ireland) should be held at ambient for a maximum of 2 hours including transit, display and service time	
	☐ Separate food safe containers must be provided for the transport of raw and ready to eat foods	
COMPASS	☐ Transport all food in suitable containers, covering or wrapping the food and transporting in a clean vehicle.	
Transportation	☐ Ensure allergen records are provided for all transported foods	
More No. 27	☐ Food above +8°C (5°C in Ireland) (for a maximum of 4 hours to be discarded)	
	☐ Food below +63°C can be reheated to +75°C (+82°C in Scotland) if not previously reheated	
	☐ At end of service discard any leftover food	
	□ Select one hot food item & one cold item per box/container and record the temperature of the food on dispatch ensuring you capture the temperature and time of dispatch	
Constitution of Annalism Constitution of Annal	☐ On arrival record the time and the temperature of food as appropriate	
	☐ Unit manager to check and sign prior to filing	
	☐ Ensure the daily vehicle safety checks are completed before departure	
HACCP Stages	Colleague Validation	More Information
Food preparationCookingFood Service	 What Temperature are colds food transported What Temperature must hot food be transported at At end of service what happens to left over foods that have been transported How many items must be checked prior to despatch? 	More information can be found within the Good Hygiene Practice Guide No: 22 Food Transportation and the HSE website

CO22_COM_POA4_030445.indd 22 03/03/2022 15:49